International Joint Commission

SAB Workshop on

An Ecosystem Approach to the

Health Effects of Mercury
in the Great Lakes Basin,

February 26/27, 2003, Windsor, Ontario


Michael Gilbertson

Presentation to the

Binational Toxics Strategy Meeting Wednesday, 14 May, 2003, Windsor, Ontario.

#### Introduction

- 1998, Health Canada published data and statistics on community health in 17 Canadian Areas of Concern;
- Selected 70+ health endpoints the "might be related to exposures to pollutants";
- Cerebral palsy included because of associatio with exposures to methylmercury;
  - Minamata, Japan, 1950s;
  - Iraq, 1960s and 1970s



#### 2001-2003 IJC Priorities

- Mercury chosen as a priority;
- Great Lakes Science Advisory Board activities
  - Consultation on the Health Effects of Mercury, Ann Arbor, February 2002;
  - Workshop on An Ecosystem Approach, Windsor, February, 2003 (IAQAB/HPTF).

### An Ecosystem Approach

- In 1978, Science Advisory Board recommendation to the International Joint Commission;
- Tool to respond to scientific complexity of persistent toxic substances:-
  - Multimedia sources and sinks;
  - Multimedia fate and distribution;
  - Multimedia routes of exposure;
  - Exposures of critical subpopulations;
  - Identify health effects and
  - Focus public health interventions and remedial actions.

### Great Lakes Science Advisory Board Consensus Workshop Summary

#### Sources, sinks, pools, exposures and effects

Milton Clark, Ph.D., U.S. EPA
Donna Mergler, Ph.D., University of Quebec
Laurie Chan, Ph.D., McGill University
Luke Trip, Senes Consulting, Ottawa
Chris DeRosa. Ph.D., ATSDR
Annette Ashizawa, Ph.D., ATSDR
Henry Anderson, M.D., WDHFS

### Anthropogenic Sources, Sinks and Pools

- Local, regional and global;
- Cycling between biosphere, earth, air and water;
- In sediment, converted to methylmercury;
- Biomagnification in food chains;
- 20% of Great Lakes loadings are from global sources and deposition;
- Need for international cooperation.

# Anthropogenic Sources, Sinks and Pools

- Legacy in Great Lakes from past practices and former industrial sites;
- Mercury cell chloralkali plants;
- Slimicide in paper manufacturing;
- Mining gold;
- Pesticide application;

# Anthropogenic Sources, Sinks and Pools

- Need further inventories for mass balance approaches to loadings, sources and fate;
- Need to reduce mercury in products and in waste streams.


### Mercury Exposure and Human Health Effects

- Mercury extensively reviewed (ATSDR 1999);
- Methylmercury extensively reviewed (NAS/NRC 2000);
  - Exposure from consumption of fish;
  - Potent neurological toxicant;
  - Mercury causes subtle, adverse neurological effects at low doses;
  - With more information, exposure level associated with adverse effects has been lowered.
- Some Great Lakes fish consumers are above the levels of potential health concerns (5.8 ppb in blood EPA).

#### **NAS/NRC Conclusion**

- Mercury Exposure and Human Health Effects
  - National Academy of Sciences (2000)
     <u>Toxicological Effects of Methyl Mercury</u>
  - Children of women who consumed large amounts of fish and seafood during pregnancy are at highest risk.
  - "...Predicted increase in the number of children who have to struggle to keep up in school and who require remedial classes or special education".

### National Trends in U.S. Fish Advisories for Various Pollutants



### Consensus Summary – Consumption Advisories

- In1990's, several Great Lakes States and the Province of Ontario issued sport fish consumption advice for women and children for all lakes;
  - For "unrestricted" category only one meal per week;

### Consensus Summary – Consumption Advisories

- In 2001, EPA issued national sportfish advice (meal per week) and
- FDA issued national advice for consumption of fish caught commercially;
- In 2003, U.S. blood survey (NHANES) 8% of women above safe levels;
  - 320,000 infants/yr at risk (EPA, 2000).

# Consensus Conclusions - Mercury Levels and Sources

- Recent declines in mercury in Great Lakes fish;
- Some AOCs and inland lake fish are still up to 10 fold higher than acceptable levels;
- Further research is needed on sources and inputs to more effectively target mercury reduction strategies.

### Health Implications and Future Research

- High intake women consumers of fish from certain AOCs or inland lakes would be expected to have mercury exposures resulting in adverse health effects in the developing child;
- Epidemiological evaluations should be undertaken in the Great Lakes area to determine if health effects exist as seen in other locations in the world.

### Risk Management and Risk Communication

- Low awareness of fish advisories among women and ethnic minorities;
- More effective risk communication approaches are needed to reduce mercury exposures;

### Risk Management and Risk Communication

- Need to balance risk communication with health benefits of fish consumption;
- Sensitivity is needed to avoid nutritional and cultural impacts (e.g., First Nations).